

**Q.1(a) Find Laplace transform of  $t e^{3t} \cos t$ . [5]**

**Ans.:**  $L\{\cos t\} = \frac{s}{s^2+1}$ ,  $L\{t \cos t\} = (-1) \frac{d}{ds} \frac{s}{s^2+1}$

$$\therefore L\{t \cos t\} = - \left\{ \frac{s^2+1-s(2s)}{(s^2+1)^2} \right\} = \frac{s^2-1}{(s^2+1)^2}$$

by F.S.T.

$$L\{t e^{3t} \cos t\} = \frac{(s-3)^2-1}{[(s-3)^2+1]^2}$$

**Q.1(b) Evaluate  $\oint_c \frac{z-1}{z^2+2z+5} dz$ , where  $c$  is  $|z+1+i|=2$ . [5]**

**Ans.:** Let  $I = \int_c \frac{z-1}{z^2+2z+5} dz$   $c : |z+1+i|=2$

for poles,  $z^2+2z+5=0$

$$\Rightarrow z = -1+2i, \quad z = -1-2i$$

$$|-1+2i+1+i| = |0+3i| = 3 > 2 \quad \text{outside}$$

$$|-1-2i+1+i| = |0-i| = 1 < 2 \quad \Rightarrow \text{pole } z = -1-2i \text{ is inside}$$

$$\begin{aligned} \text{at } z = -1-2i \quad \text{Residue} &= \frac{z-1}{2z+2} \Big|_{z=-1-2i} = \frac{-1-2i-1}{-2-4i+2} \\ &= \frac{-2-2i}{-4i} = \frac{1+i}{2i} \end{aligned}$$

$$\text{by Residue Theorem } \int_c \frac{z-1}{z^2+2z+5} = 2\pi i \left( \frac{1+i}{2i} \right) = \pi(1+i)$$

**Q.1(c) Show that  $f(z) = \sinh z$  is analytic. Hence find its derivative. [5]**

**Ans.:**  $f(z) = \sinh z = \sinh(x+iy) = \sinh x \cosh iy + \cosh x \sinh iy$   
 $u+iv = \sinh x \cos y + i \cosh x \sin y$

$$\Rightarrow u = \sinh x \cos y$$

$$u_x = \cosh x \cos y \quad \dots(1)$$

$$u_y = \sinh x \sin y \quad \dots(2)$$

$$v = \cosh x \sin y$$

$$v_x = \sinh x \sin y \quad \dots(3)$$

$$v_y = \cosh x \cos y \quad \dots(4)$$

From (1), (4) and (2), (3)  $u_x = v_y$ ,  $v_x = -u_y$

$\therefore f(z) = \sinh z$  is analytic.

$$\text{Now } f'(z) = u_x + iv_x \Big|_{x=z, y=0} = \cosh x \cos y + i \sinh x \sin y \Big|_{x=z, y=0}$$

$$f'(z) = \cosh z$$

**Q.1(d) Compute spearman’s rank correlation for the data :** [5]

<b>X :</b>	<b>18</b>	<b>20</b>	<b>34</b>	<b>52</b>	<b>12</b>
<b>Y :</b>	<b>39</b>	<b>23</b>	<b>35</b>	<b>18</b>	<b>46</b>

**Ans.:**

X	Y	R <sub>x</sub>	R <sub>y</sub>	d = R <sub>x</sub> – R <sub>y</sub>	d <sup>2</sup>
18	39	4	2	2	4
20	23	3	4	-1	1
34	35	2	3	-1	1
52	18	1	5	-4	16
12	46	5	1	4	16

$$\sum d^2 = 38$$

$$R = 1 - \frac{6 \sum d^2}{n(n^2 - 1)} = 1 - \frac{6(38)}{5(24)} \Rightarrow R = -0.9$$

**Q.2(a) Show that the function  $\omega = \frac{4}{z}$  transforms the straight line  $x = c$  in the  $z$ -plane into circle in  $w$ -plane. Find its centre and radius.** [6]

**Ans.:**  $x = c$  ... (1)

Given  $\omega = \frac{4}{z} \Rightarrow z = \frac{4}{\omega} \Rightarrow x + iy = \frac{4}{u + iv}$

$\therefore x + iy = \frac{4u}{u^2 + v^2} - \frac{i4v}{u^2 + v^2} \Rightarrow x = \frac{4u}{u^2 + v^2}$

using this in (1)  $\frac{4u}{u^2 + v^2} = c \Rightarrow u^2 + v^2 - \frac{4u}{c} = 0$

centre =  $\left(\frac{z}{c}, 0\right)$ , rad =  $\frac{2}{c}$

**Q.2(b) Show that  $\int_0^\infty e^{-t} \int_0^t \frac{\sin u}{u} du dt = \frac{\pi}{4}$**  [6]

**Ans.:**  $\int_0^\infty e^{-t} \int_0^t \frac{\sin u}{u} du dt = L \left\{ \int_0^t \frac{\sin u}{u} du \right\} \Big|_{s=1}$

Now

$$\begin{aligned} L \left\{ \int_0^t \frac{\sin u}{u} du \right\} &= \frac{1}{s} L \left\{ \frac{\sin u}{u} \right\} = \frac{1}{s} \int_s^\infty \frac{1}{s^2 + 1} ds \\ &= \frac{1}{s} \left\{ \tan^{-1} s \right\}_{s=s}^{s=\infty} = \frac{1}{s} \left[ \frac{\pi}{2} - \tan^{-1} s \right] \end{aligned}$$

$\therefore$  at  $s = 1$ ,

$$= \frac{1}{1} \left[ \frac{\pi}{2} - \tan^{-1} 1 \right] = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}$$

**Q.2(c)** Obtain fourier series for  $f(x) = \begin{cases} 1 + \frac{2x}{\pi} & -\pi \leq x \leq 0 \\ 1 - \frac{2x}{\pi} & 0 \leq x \leq \pi \end{cases}$  [8]

Hence deduce that  $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

**Ans.:**  $f(x)$  is even function.  $\Rightarrow b_n = 0$

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos(nx) \quad \dots(1)$$

$$a_0 = \frac{1}{2\pi} 2 \int_0^{\pi} \left(1 - \frac{2x}{\pi}\right) dx = \frac{1}{\pi} \left[ x - \frac{x^2}{\pi} \right]_0^{\pi} \Rightarrow a_0 = 0$$

$$a_n = \frac{1}{\pi} 2 \int_0^{\pi} \left(1 - \frac{2x}{\pi}\right) \cos(nx) dx = \frac{2}{\pi} \left\{ \left(1 - \frac{2x}{\pi}\right) \left(\frac{\sin(nx)}{n}\right) - \left(-\frac{2}{\pi}\right) \left(\frac{-\cos(nx)}{n^2}\right) \right\}_0^{\pi}$$

$$= \frac{2}{\pi} \left\{ \left[ 0 - \frac{2}{\pi n^2} (-1)^n \right] - \left[ 0 - \frac{2}{\pi n^2} \right] \right\}$$

$$a_n = \frac{4}{n^2 \pi^2} [1 - (-1)^n]$$

using this in (1)

$$f(x) = \sum_{n=1}^{\infty} \frac{4}{n^2 \pi^2} [1 - (-1)^n] \cos(nx)$$

for deduction put  $x = 0$  and note that  $f(0) = 1$

$$1 = \sum_{n=1}^{\infty} \frac{4}{\pi^2 n^2} [1 - (-1)^n]$$

$$1 = \frac{4}{\pi^2} \left[ \frac{2}{1^2} + 0 + \frac{2}{3^2} + 0 + \frac{2}{5^2} + \dots \right] \Rightarrow \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

**Q.3(a)** Evaluate  $\oint_c \frac{e^{kz}}{z} dz$ , where  $c: |z| = 1$ . [6]

Hence deduce that  $\int_0^{\pi} e^{k \sin \theta} \cos(k \sin \theta) d\theta = \pi$

**Ans.:**  $I = \int_c \frac{e^{kz}}{z} dz$ ,  $c: |z| = 1$

pole  $z = 0$  is inside  $|z| = 1$

$$\therefore \int_c \frac{e^{kz}}{z} dz = 2\pi i f(0), \quad \text{where } f(z) = e^{kz}$$

$$= 2\pi i \quad (1)$$

put  $z = e^{i\theta} \Rightarrow dz = ie^{i\theta} d\theta$ , limits 0 to  $2\pi$

$$\therefore (1) \Rightarrow \int_0^{2\pi} \frac{e^{ke^{i\theta}}}{e^{i\theta}} i e^{i\theta} d\theta = 2\pi i$$

$$\begin{aligned}
 i \int_0^{2\pi} e^{k(\cos \theta + i \sin \theta)} d\theta &= 2\pi i \\
 \int_0^{2\pi} e^{k \cos \theta} [\cos(k \sin \theta) + i \sin(k \sin \theta)] d\theta &= 2\pi \\
 \int_0^{2\pi} e^{k \cos \theta} \cos(k \sin \theta) d\theta + \int_0^{2\pi} e^{k \cos \theta} i \sin(k \sin \theta) d\theta &= 2\pi + 0i \\
 \Rightarrow \int_0^{2\pi} e^{k \cos \theta} \cos(k \sin \theta) d\theta &= 2\pi \\
 2 \int_0^{2\pi} e^{k \cos \theta} \cos(k \sin \theta) d\theta &= 2\pi \\
 \Rightarrow \int_0^{2\pi} e^{k \cos \theta} \cos(k \sin \theta) d\theta &= \pi
 \end{aligned}$$

**Q.3(b)** For the lines of regression  $6y - 5x = 90$ ,  $15x - 8y = 130$  and  $\sigma_x^2 = 16$  [6]

Find (i)  $\bar{x}$ ,  $\bar{y}$  (ii)  $r$  (iii)  $\sigma_y$

**Ans.:** given  $6y - 5x = 90$ ,  $15x - 8y = 130$   
 solving we get  $x = 30$ ,  $y = 40$   
 i.e.  $\bar{x} = 30$ ,  $\bar{y} = 40$

$$y = \frac{5}{6}x + 15 \Rightarrow b_{yx} = \frac{5}{6}$$

$$x = \frac{8}{15}y + \frac{130}{15} \Rightarrow b_{xy} = \frac{8}{15}$$

$$\therefore r = \pm \sqrt{b_{xy} b_{yx}} = +\sqrt{\frac{5}{6} \cdot \frac{8}{15}} \Rightarrow r = 0.67$$

$$\text{given } \sigma_x^2 = 16 \Rightarrow \sigma_x = 4$$

$$b_{xy} = \frac{8}{15} \Rightarrow r \frac{\sigma_x}{\sigma_y} = \frac{8}{15} \Rightarrow \frac{2\left(\frac{4}{\sigma_y}\right)}{3} = \frac{8}{15}$$

$$6\sigma_y = 5$$

**Q.3(c)** Solve the differential equation  $\frac{dy}{dx} + 2y + \int_0^t y dt = \sin t$  using Laplace [8]

transform give  $y(0) = 1$

$$\text{Ans.} \quad L\{y'(t)\} + 2L\{y(t)\} + L\left\{\int_0^t y(t) dt\right\} = L\{\sin t\}$$

$$5y(s) - 1 + 2y(s) + \frac{1}{s}y(s) = \frac{1}{s^2 + 1}$$

$$\left(s + 2 + \frac{1}{s}\right)y(s) = \frac{1}{s^2 + 1} + 1 = \frac{s^2 + 2}{s^2 + 1}$$

$$\frac{(s^2 + 2s + 1)}{s} y(s) = \frac{s^2 + 2}{s^2 + 1} \Rightarrow y(s) = \frac{s(s^2 + 2)}{(s+1)^2 (s^2 + 1)}$$

$$y(s) = \frac{a}{s+1} + \frac{b}{(s+1)^2} + \frac{cs+d}{s^2+1}, a=1, b=-\frac{3}{2}, c=0, d=\frac{1}{2}$$

$$\therefore y(s) = \frac{1}{s+1} - \frac{3}{2} \frac{1}{(s+1)^2} + \frac{1}{2} \frac{1}{s^2+1}$$

taking inverse Laplace,

$$y(t) = e^{-t} - \frac{3}{2} te^{-t} + \frac{1}{2} \sin t$$

**Q.4(a) Find Laurent's series for  $f(z) = \frac{2}{(z-1)(z-2)}$  indicating region of convergence. [6]**

**Ans.:**  $f(z) = \frac{2}{(z-1)(z-2)} = \frac{a}{z-1} + \frac{b}{z-2} = -\frac{2}{z-1} + \frac{2}{z-2} \dots(1)$

for Laurent's series ROC is (1)  $|z| > 1, |z| > 2$  (ii)  $1 < |z| < 2$

case (1):  $|z| > 1, |z| > 2$

$$(1) \Rightarrow f(z) = \frac{-2}{z\left(1-\frac{1}{z}\right)} + \frac{2}{z\left(1-\frac{2}{z}\right)} = -\frac{2}{z}\left(1-\frac{1}{z}\right)^{-1} + \frac{2}{z}\left(1-\frac{2}{z}\right)^{-1}$$

$$f(z) = \frac{-2}{z}\left[1 + \frac{1}{z} + \frac{1}{z^2} + \dots\right] + \frac{2}{z}\left[1 + \frac{2}{z} + \frac{4}{z^2} + \dots\right]$$

case (2):  $1 < |z| < 2$

$$(1) \Rightarrow f(z) = \frac{-2}{z\left(1-\frac{1}{z}\right)} - \frac{2}{2\left(1-\frac{z}{2}\right)} = -\frac{2}{z}\left(1-\frac{1}{z}\right)^{-1} - \left(1-\frac{z}{2}\right)^{-1}$$

$$f(z) = \frac{-2}{z}\left[1 + \frac{1}{z} + \frac{1}{z^2} + \dots\right] - \left[1 + \frac{z}{2} + \frac{z^2}{4} + \dots\right]$$

**Q.4(b) Show that  $\cos x = 8\pi \sum_{m=1}^{\infty} \frac{m}{4m^2-1} \sin(2mx)$ , if  $0 < x < \pi$  [6]**

**Ans.:** Half range sine series  $f(x) = \sum_{n=1}^{\infty} b_n \sin(nx) \dots(1)$

$$b_n = \frac{2}{\pi} \int_0^{\pi} \cos x \sin(nx) dx = \frac{2}{\pi} \int_0^{\pi} \left[ \frac{\sin(nx+x)}{2} + \frac{\sin(nx-x)}{2} \right] dx$$

$$= \frac{1}{\pi} \left[ \frac{-\cos(nx+x)}{n+1} - \frac{\cos(nx-x)}{n-1} \right]_0^{\pi} = \frac{1}{\pi} \left[ \frac{(-1)^n}{n+1} + \frac{(-1)^n}{n-1} \right] - \left[ -\frac{1}{n+1} - \frac{1}{n-1} \right]$$

$$b_n = \frac{1}{\pi} [1 + (-1)^n] \left[ \frac{1}{n+1} + \frac{1}{n-1} \right] = \frac{2n}{\pi(n^2-1)} [1 + (-1)^n]$$

$n \neq 1$

$$\begin{aligned} \text{Now } b_1 &= \frac{2}{\pi} \int_0^{\pi} \sin x \cos x \, dx = \frac{1}{\pi} \int_0^{\pi} \sin 2x \, dx \\ &= \frac{1}{\pi} \left[ -\frac{\cos 2x}{2} \right]_0^{\pi} = -\frac{1}{2\pi} [1-1] \Rightarrow b_1 = 0 \end{aligned}$$

$$\Rightarrow \cos x = \sum_{n=2}^{\infty} \frac{2n}{\pi(n^2-1)} [1+(-1)^n] \sin(nx)$$

put  $n = 2m$

$$\therefore \cos x = \sum_{2m=2}^{\infty} \frac{2(2m)}{\pi(4m^2-1)} [1+(-1)^{2m}] \sin(2mx)$$

$$\therefore \cos x = \frac{8}{\pi} \sum_{m=1}^{\infty} \frac{m}{4m^2-1} \sin(2mx)$$

**Q.4(c) Find Bilinear transformation which maps the points 1, i, -1 onto the points i, 0, -i. Hence find fixed points and image of  $|z| < 1$ . [8]**

**Ans.:** Consider B. l.  $\omega = \frac{az+b}{cz+d}$  ... (1)

given  $z = 1, \omega = i$  (1)

$$\Rightarrow i = \frac{a+b}{c+d} \Rightarrow ic + id = a + b \quad \dots(2)$$

given  $z = i, \omega = 0$  (1)

$$\Rightarrow 0 = \frac{ia+b}{ic+d} \Rightarrow b = -ia \quad \dots(3)$$

given  $z = -1, \omega = -i$  (1)

$$\Rightarrow -i = \frac{-a+b}{-c+d} \Rightarrow ic - id = -a + b \quad \dots(4)$$

$$(2) + (4) \Rightarrow 2ic = 2b \Rightarrow 2ic = -2ia \Rightarrow c = -a \quad \dots(5)$$

$$(2) - (4) \Rightarrow 2id = 2a \Rightarrow d = -ia \quad \dots(6)$$

using (3), (5), (6), in (1)

$$\omega = \frac{az-ia}{-az-ia} \Rightarrow \omega = \frac{i-z}{i+z} \quad \dots(7)$$

For fixed points  $w = z$ , (7)  $\Rightarrow z = \frac{i-z}{i+z}$

$$z^2 + iz = i - z \Rightarrow z^2 + (i+1)z - i = 0$$

$$z = \frac{-i-1 \pm \sqrt{-1+1+2i+4i}}{2} = \frac{-i-1 \pm \sqrt{6i}}{2}$$

to find image of  $|z| < 1$  : (7)  $\Rightarrow i\omega + \omega z = i - z \Rightarrow (1 + \omega)z = i - i\omega$

$$z = \frac{i - i(u+iv)}{1+u+iv}$$

$$\therefore |z| < 1 \Rightarrow \left| \frac{i - iu + v}{1 + u + iv} \right| < 1$$

$$|i - iu + v| = |1 + u + iv|$$

$$\begin{aligned} \sqrt{v^2 + (1-u)^2} &< \sqrt{(1+u)^2 + v^2} \\ \Rightarrow v^2 + 1 - 2u + u^2 &< 1 + 2u + u^2 + v^2 \Rightarrow 0 < 4u \\ \Rightarrow 0 < u \end{aligned}$$

**Q.5(a) Solve using Bender-Schmidt method  $\frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t} = 0$ , subject to the conditions [6]**

**$u(0, t) = 0, u(1, t) = 0, u(x, 0) = \sin \pi x, 0 \leq x \leq 1$**

**Ans.:**  $a = 1$ , take  $h = 0.2 \Rightarrow x : 0, 0.2, 0.4, 0.6, 0.8, 1$


$$k = \frac{a}{2} h^2 = 0.02$$

$t : 0, 0.02, 0.04, 0.06, 0.08, 0.10$   
 $x \rightarrow h = 0.2$

t ↓ k = 0.02	t \ x	0.0	0.2	0.4	0.6	0.8	1
	0.00	0	0.5878	0.9511	0.9511	0.5878	0
0.02	0						0
0.04	0						0
0.06	0						0
0.08	0						0
0.10	0						0

using Bender-Schmidt formula  $c = \frac{(a+b)}{2}$

t \ x	0	0.2	0.4	0.6	0.8	1
0	0	0.5878	0.9511	0.9511	0.5878	0
0.02	0	0.4756	0.7695	0.7695	0.4756	0
0.04	0	0.3848	0.6225	0.6225	0.3848	0
0.06	0	0.3113	0.5036	0.5036	0.3113	0
0.08	0	0.2518	0.4074	0.4074	0.2518	0
0.10	0	0.2037	0.3296	0.3296	0.2037	0



**Q.5(b) Determine the solution of one-dimensional heat equation under the [6]**

**boundary conditions,  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$**

**$u(0,t) = 0, u(\ell, t) = 0, u(x,0) = x, (0 < x < \ell), \ell$  being length of the rod.**

**Ans.:** Solution of heat flow is given by

$$u = (c_1 \cos mx + c_2 \sin mx) e^{-m^2 c^2 t} \quad \dots(1)$$

given  $u = 0$  when  $x = 0$

$$(1) \Rightarrow 0 = c_1 e^{-m^2 c^2 t} \Rightarrow c_1 = 0$$

$$\therefore (1) \Rightarrow u = c_2 \sin mx e^{-m^2 c^2 t} \quad \dots(2)$$

given when  $x = \ell, u = 0$

$$(2) \Rightarrow 0 = c \sin m \ell e^{-m^2 c^2 t} \Rightarrow \sin m \ell = 0 \Rightarrow m \ell = n\pi$$

$$\Rightarrow m = \frac{n\pi}{\ell}$$

$$(2) \Rightarrow u = c_2 \sin\left(\frac{n\pi x}{\ell}\right) e^{-\frac{n^2 \pi^2}{\ell^2} c^2 t} \quad \dots(3)$$

adding all above solutions for  $n = 1, 2, \dots$  we get the general solution

$$u = \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{\ell}\right) e^{-\frac{n^2 \pi^2 c^2 t}{\ell^2}} \quad \dots(3)$$

given when  $t = 0, u = x$

$$(3) \Rightarrow x = \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{\ell}\right)$$

which is half range series in  $(0, \ell)$

$$b_n = \frac{2}{\ell} \int_0^{\ell} x \sin\left(\frac{n\pi x}{\ell}\right) dx = \frac{2}{\ell} \left\{ (x) \left[ \frac{-\cos\left(\frac{n\pi x}{\ell}\right)}{\frac{n\pi}{\ell}} \right] - (1) \left[ \frac{-\sin\left(\frac{n\pi x}{\ell}\right)}{\frac{n^2 \pi^2}{\ell^2}} \right] \right\}_0^{\ell}$$

$$= \frac{2}{\ell} \left\{ \left[ -\frac{\ell}{n\pi} (-1)^n - 0 \right] - [0 - 0] \right\} = -\frac{2\ell}{n\pi} (-1)^n$$

using this in (3) we get,

$$u = \sum_{n=1}^{\infty} \frac{-2\ell}{n\pi} (-1)^n \sin\left(\frac{n\pi x}{\ell}\right) e^{-\frac{n^2 \pi^2 c^2 t}{\ell^2}}$$

**Q.5(c) Find inverse Laplace Transform of**

**[8]**

(i)  $\log\left(1 + \frac{a^2}{s^2}\right)$       (ii)  $\frac{e^{-s}}{s^2 + s + 1}$

**Ans.:** (i) Let  $f(s) = \log\left(1 + \frac{a^2}{s^2}\right) = \log\left(\frac{s^2 + a^2}{s^2}\right)$

$$f(s) = \log(s^2 + a^2) - \log s^2$$

$$\Rightarrow f'(s) = \frac{2s}{s^2 + a^2} - \frac{2}{s}$$

taking  $L^{-1}$  we get

$$L^{-1}\{f'(s)\} = 2 \cos at - 2$$

$$\Rightarrow -t f(t) = 2 \cos at - 2$$

$$\Rightarrow f(t) = \frac{2(1 - \cos at)}{t}$$



$$(ii) \text{ Let } f(s) = \frac{1}{s^2 + s + 1} = \frac{1}{\left(s + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$\therefore L^{-1}\{F(s)\} = L^{-1}\left\{\frac{1}{\left(s + \frac{1}{2}\right)^2 + \frac{3}{4}}\right\}$$

$$\Rightarrow F(t) = e^{-\frac{1}{2}t} L^{-1}\left\{\frac{1}{s^2 + \frac{3}{4}}\right\} = \frac{e^{-\frac{1}{2}t}}{\frac{\sqrt{3}}{2}} \sin\left(\frac{\sqrt{3}}{2}t\right)$$

we know that,

$$L^{-1}\{e^{-as} f(s)\} = f(t-a) H(t-a)$$

$$\therefore L^{-1}\left\{\frac{e^{-s}}{s^2 + s + 1}\right\} = \frac{2}{\sqrt{3}} e^{-\frac{1}{2}(t-1)} \sin\left[\frac{\sqrt{3}}{2}(t-1)\right] \cdot H(t-1)$$

**Q.6(a) Obtain complex form of fourier series for  $F(x) = e^{\alpha x}$ , in  $(-\pi, \pi)$  where  $a$  is not an integer. [6]**

**Ans.:** Complex form of fourier series is given by

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{inx} \quad \dots(1)$$

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ax} e^{-inx} dx = \frac{1}{2\pi} \left\{ \frac{e^{ax-inx}}{a-in} \right\}_{-\pi}^{\pi} = \frac{1}{2\pi} \frac{(a+in)}{(a+in)(a-in)} \{e^{a\pi} e^{-in\pi} - e^{-a\pi} e^{in\pi}\}$$

$$= \frac{1}{2\pi} \frac{(a+in)}{(a^2+n^2)} (-1)^n (e^{a\pi} - e^{-a\pi}) = \frac{(a+in)}{\pi(a^2+n^2)} (-1)^n \sinh(a\pi)$$

using this in (1)

$$e^{ax} = \sum_{n=-\infty}^{\infty} \frac{(a+in)}{\pi(a^2+n^2)} (-1)^n \sinh(a\pi) e^{inx}$$

**Q.6(b) Fit a curve  $y = a \cdot b^x$  to the following data, using method of least squares. [6]**

<b>X :</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
<b>Y :</b>	<b>144</b>	<b>172.8</b>	<b>207.4</b>	<b>248.8</b>	<b>298.5</b>

**Ans.:**  $y = a \cdot b^x$  (taking log) ... (1)

$$\log y = \log a + x \log b$$

$$\Rightarrow \sum \log y = n \ln a + \ln b \sum x \quad \dots(2)$$

$$\sum x \log y = \ln a \sum x + \ln b \sum x^2 \quad \dots(3)$$

$$\left. \begin{aligned} (2) &\Rightarrow 26.58 = 5A + 20B \\ (3) &\Rightarrow 108.51 = 20A + 90B \end{aligned} \right\} \text{ where } A = \ln a, B = \ln b$$

$$\text{Solving} \quad \ln a = 4.62$$

$$B = 0.179 \quad a = 101.49$$

$$\therefore (1) \Rightarrow \ln b = 0.179$$

$$y = (101.49) (1.196)^x \quad b = 1.196$$

Q.6(c) (i) Evaluate  $\int_0^{2\pi} \frac{d\theta}{5 + 3 \sin \theta}$  using Residue Theorem. [4]

(ii) Evaluate  $\int_{-\infty}^{\infty} \frac{dx}{x^2 + 1}$  using Residue Theorem. [4]

Ans.: (i)  $I = \int_0^{2\pi} \frac{d\theta}{5 + 3 \sin \theta} = \text{put } z = e^{i\theta} \Rightarrow d\theta = \frac{dz}{iz}$

$$\sin \theta = \frac{z^2 - 1}{2iz}$$

$$\therefore I = \int_c \frac{\frac{dz}{iz}}{5 + 3 \left( \frac{z^2 - 1}{2i} \right)} = \int_c \frac{2}{3z^2 + 10iz - 3} dz$$

poles are  $z = -\frac{i}{3}, z = -3i$

$z = -\frac{i}{3}$  lies inside  $|z| = 1$

at  $z = -\frac{i}{3}$ ,  $\text{Res} = \frac{2}{6z + 10i} \Big|_{z = -\frac{i}{3}} = \frac{2}{-2i + 10i} = \frac{1}{4i} = \frac{1}{4i}$

by Residue Theorem,

$$I = 2\pi i \left( \frac{1}{4i} \right) = \frac{\pi}{2}$$

(ii) Consider  $\int_c \frac{dz}{z^2 + 1}$  where  $c$  is large semicircle as shown in figure.

$$\therefore \int_{-R}^R \frac{dz}{z^2 + 1} + \int_{C_1} \frac{dz}{z^2 + 1} = \int_c \frac{dz}{z^2 + 1} \quad \dots(1)$$

taking limit as  $R \rightarrow \infty$

along real axis  $y = 0 \Rightarrow z = x$  and  $\int_{C_1} \frac{dz}{z^2 + 1} = 0$

$$\therefore (1) \Rightarrow \int_{-\infty}^{\infty} \frac{dx}{x^2 + 1} = \int_c \frac{dz}{z^2 + 1},$$

poles are  $z = \pm i$  but  $z = i$  lies inside

at  $z = i$   $\text{Res} = \frac{1}{2z} \Big|_{z=i} = \frac{1}{2i}$

$\therefore$  by residue theorem

$$\int_{-\infty}^{\infty} \frac{dx}{x^2 + 1} = 2\pi i \left( \frac{1}{2i} \right) = \pi$$

